

Here for

More



Caring for your eyes while taking ELAHERE

Use this brochure to learn about what you can do to help manage the risk of eye-related side effects during treatment with ELAHERE.

ELAHERE is a prescription medicine used to treat adults with folate receptor-alpha (FR α) positive ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who:

- have not responded to or are no longer responding to treatment with platinum-based chemotherapy **and**
- have received 1 to 3 prior types of chemotherapy.

Your healthcare provider will perform a test to make sure that ELAHERE is right for you.

It is not known if ELAHERE is safe and effective in children.

Actor portrayals.

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.

Continue reading information about eye problems on pages 8–10.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

What information to expect within this guide



- Important Safety Information3
- How could ELAHERE affect my eyes?.....8
- What should I know before starting ELAHERE?9
- Monitor your eyes for side effects.....10
- Recommended schedule for eye drops 12
- Tips to help address vision changes 13
- Talk to your doctor about eye care while on ELAHERE..... 16

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

USE and IMPORTANT SAFETY INFORMATION



What is ELAHERE?

ELAHERE is a prescription medicine used to treat adults with folate receptor-alpha positive ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who:

- have not responded to or are no longer responding to treatment with platinum-based chemotherapy **and**
- have received 1 to 3 prior types of chemotherapy.

Your healthcare provider will perform a test to make sure that ELAHERE is right for you.

It is not known if ELAHERE is safe and effective in children.

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.

- Your healthcare provider will send you to see an eye care professional to check your eyes before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed for any worsening signs and symptoms of eye problems.
- Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before you start and during your treatment with ELAHERE. You should use eye drops as directed by your healthcare provider.
- Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (cont'd)



What should I tell my healthcare provider before receiving ELAHERE?

Tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems.
- have numbness or tingling in your hands or feet.
- have liver problems.
- are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.

Patients who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ELAHERE.
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.
- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (cont'd)



What are the possible side effects of ELAHERE?

ELAHERE can cause serious side effects, including:

- **Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.
- **Lung problems (pneumonitis).** ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain.
- **Peripheral neuropathy.** Nerve problems called peripheral neuropathy are common during treatment with ELAHERE and can also be severe. Your healthcare provider will monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if you get new or worsening numbness, tingling, burning sensation or pain in your hands or feet or muscle weakness.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (cont'd)



The most common side effects and abnormal labs of ELAHERE include:

- increased liver enzymes in the blood
- feeling tired
- blurred vision
- nausea
- diarrhea
- stomach-area (abdominal) pain
- changes in the cornea (part of the eye)
- peripheral neuropathy
- muscle, bone, or joint pain
- decreased red or white blood cell counts
- decreased platelets
- decreased magnesium level in the blood
- dry eye
- constipation
- vomiting
- decreased albumin level in the blood
- decreased appetite

Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have certain side effects.

These are not all of the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).



Actor portrayals.

How could ELAHERE affect my eyes?



People treated with ELAHERE can develop eye-related side effects, which can be severe. These may include blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes. Tell your doctor if you have a history of vision or eye problems.



Eye-related side effects most commonly occurred between the second and third infusion of ELAHERE



1% of people in the study stopped treatment because of eye problems

Avoid wearing contact lenses throughout your treatment with ELAHERE unless your doctor tells you that you can. **Your doctor will also have you meet with an eye doctor (optometrist or ophthalmologist) before your first infusion and periodically throughout your treatment. You will receive instructions for eye drops to use during your treatment. Be sure to use them as directed.**

Explore the Eye Drop Tracker

While on treatment, use this tool to help track your eye drop use, notes to share with your doctor, and upcoming appointments

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

What should I know before starting ELAHERE?



Before starting treatment, tell your oncologist and eye doctor about any eye problems you may have



You will visit an eye doctor for a baseline eye exam. You will then have follow-up eye exams throughout your treatment on ELAHERE, about every 6 weeks during the first 8 cycles of treatment. This will help your healthcare team keep track of any potential changes to your vision or eyes during treatment.



Your doctor may recommend 2 different types of eye drops to help reduce the risk of developing eye-related side effects:

- Prescription steroid eye drops, filled by a pharmacist
- Preservative-free lubricating eye drops, which can be purchased over the counter

Apply the eye drops as directed by your doctor.

There may be additional costs associated with eye exams. Please check with your insurance provider for more information

IMPORTANT SAFETY INFORMATION (CONT'D)

- Your healthcare provider will send you to see an eye care professional to check your eyes before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed for any worsening signs and symptoms of eye problems.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

Monitor your eyes for side effects



Report any eye changes or problems to your eye doctor

You may experience some of the following eye-related side effects. Your eye doctor may be able to help you manage side effects, regardless of severity. Some eye changes may include:

- Blurred vision
- Eye pain
- Vision impairment/loss
- Dry eyes
- Eye redness
- Temporary changes to the surface of the eye
- Light sensitivity
- Cataract

If you experience side effects, your doctors may talk with each other about next steps such as:

- Your oncologist may pause treatment until your symptoms improve. Once symptoms improve, you may resume treatment
- Your oncologist may decide to continue treatment with ELAHERE, but at a reduced dose
- Depending on how serious the eye-related side effects are, your oncologist may have you discontinue treatment

IMPORTANT SAFETY INFORMATION (CONT'D)

- Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before you start and during your treatment with ELAHERE. You should use eye drops as directed by your healthcare provider.
- Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

Actor portrayals.



Recommended schedule for eye drops




This eye drop schedule shows how often you should use steroid eye drops as directed by your doctor, and how this will change as you progress through your infusion cycles with ELAHERE.

Steroid eye drops

Using steroid eye drops is recommended while taking ELAHERE.

Steroid Eye Drop Schedule for Each ELAHERE Infusion Cycle (Every 3 Weeks)



Lubricating eye drops

The use of preservative-free lubricating eye drops is also recommended at least 4 times daily and as needed during treatment. **Wait at least 10 minutes after applying steroid eye drops** before using lubricating eye drops.

Need reminders to use your eye drops? There are several mobile apps available for download, designed to help you and/or your caregiver remember your eye drop schedule. You or your caregiver can find a variety to choose from at your preferred app store

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

Tips to help address vision changes



If you experience blurred vision or vision changes, you should contact your doctor

On days when your vision is blurry, you may need to make the type bigger on your devices. Here are some ways to do so (specific examples based on your phone model may vary):



iPhone

Go to Settings, then Accessibility, then select Display & Text Size. Tap Larger Text for larger font options. Drag the slider to the size you want.

Android

Open your device's Settings app. Select Accessibility, then Text and Display. Select Font Size, and use the slider to select the size you want.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider before receiving ELAHERE?

Tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems.
- have numbness or tingling in your hands or feet.
- have liver problems.
- are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.

Patients who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ELAHERE.
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

Tips to help address vision changes (cont'd)



Computer

Did you know you can increase the font size on a website you're viewing? For example, if you have a mouse with a wheel, on a PC you can hold the Ctrl key on the keyboard and then scroll the wheel down to increase text size (scroll up to decrease text size). Even without a mouse, key commands can do the same thing. Search online for tips specific to your device.



As you will need to avoid wearing contact lenses during your treatment with ELAHERE, you may find it helpful to wear reading glasses, or to buy a different strength at your local pharmacy or other retail stores. Talk to your eye doctor about this and other eyewear options.

IMPORTANT SAFETY INFORMATION (CONT'D)

- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.

What are the possible side effects of ELAHERE?

ELAHERE can cause serious side effects, including:

- **Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.
- **Lung problems (pneumonitis).** ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including [Boxed WARNING](#) and [Medication Guide](#).



Actor portrayals.

Talk to your doctor about eye care while on ELAHERE



See below for questions to ask your doctors about eye-related side effects during treatment with ELAHERE

- What are the eye-related side effects that can occur with ELAHERE?
- Is there any way I can manage eye-related side effects during treatment?
- When should I let you know if I experience anything new or unusual with my eyes?
- Is there anything I should do to take care of my eyes?
- How often should I have an eye exam?
- **Question for your eye doctor:** My oncologist put me on a drug that can affect my eyes. Would your office be able to work with my oncologist to help monitor my eye health during my treatment?

Find an eye doctor near you.

Before starting ELAHERE treatment, you will need to meet with an eye doctor and have an eye exam

IMPORTANT SAFETY INFORMATION (CONT'D)

- **Peripheral neuropathy.** Nerve problems called peripheral neuropathy are common during treatment with ELAHERE and can also be severe. Your healthcare provider will monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if you get new or worsening numbness, tingling, burning sensation or pain in your hands or feet or muscle weakness.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

Actor portrayal.

Here for

More



Talk to your doctor if you have any questions about eye care throughout your treatment journey

To learn more and find additional resources, visit [ELAHERE.com](https://www.elaHERE.com)

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.

Continue reading information about eye problems on pages 8–10.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

immunogen

© 2024 ImmunoGen, Inc., All Rights Reserved. US-MIRV-2300320 03/2024
ImmunoGen, ELAHERE, and their respective logos are trademarks or registered trademarks of ImmunoGen, Inc.