



Actor portrayals.

# Here for

# More



ELAHERE is a prescription medicine used to treat adults with folate receptor-alpha (FR $\alpha$ ) positive ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who:

- have not responded to or are no longer responding to treatment with platinum-based chemotherapy **and**
- have received 1 to 3 prior types of chemotherapy.

Your healthcare provider will perform a test to make sure that ELAHERE is right for you.

It is not known if ELAHERE is safe and effective in children.

## SELECT IMPORTANT SAFETY INFORMATION

**What is the most important information I should know about ELAHERE?**

**ELAHERE can cause serious side effects, including:**

**Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes. **Continue reading information about eye problems on pages 3 and 5.**

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**



Actor portrayals.

# What information to expect within this guide



This brochure is designed to help you understand what to expect while on treatment with ELAHERE and support that is available.

- Understanding platinum-resistant ovarian cancer.....5
- The role of FR $\alpha$ .....6
- How ELAHERE works.....7
- How ELAHERE may help.....8
- Side effects with ELAHERE.....12
- Eye care during treatment .....15
- Before starting ELAHERE .....19
- ELAHERE dosing and infusion schedule.....21
- Questions to ask your doctor.....23
- Support services.....25

FR $\alpha$ =folate receptor alpha.

## IMPORTANT SAFETY INFORMATION (CONT'D)

- Your healthcare provider will send you to see an eye care professional to check your eyes before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed for any worsening signs and symptoms of eye problems.
- Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before you start and during your treatment with ELAHERE. You should use eye drops as directed by your healthcare provider.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including [Boxed WARNING](#) and [Medication Guide](#).**



Actor portrayals.

# A targeted option is here when it's time to consider a change in treatment



## After ovarian cancer becomes platinum resistant, what's next?

- If you or your loved one has platinum-resistant ovarian cancer, this means that the cancer came back or progressed after certain types of chemotherapy called platinum-based chemotherapy, which contains a material called platinum
- When platinum-based chemotherapy no longer works, a change in treatment approach may be needed
- If your cancer is positive for a kind of protein called folate receptor alpha, or FR $\alpha$  (FR $\alpha$ ), you may be eligible for targeted treatment with ELAHERE
- Speak to your doctor to learn more about ELAHERE and if it may be right for you

## Is ELAHERE right for me?



Your ovarian cancer is platinum resistant



Your cancer is positive for a type of protein called FR $\alpha$



You have received 1 to 3 prior types of chemotherapy



Throughout this brochure, the term “ovarian cancer” is used to refer to ovarian cancer, fallopian tube cancer, and primary peritoneal cancer

## IMPORTANT SAFETY INFORMATION (CONT'D)

- Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

# What is FR $\alpha$ and why is it important?



**About 1 in 3 women** with platinum-resistant ovarian cancer will have high levels of FR $\alpha$

**FR $\alpha$  is a protein** found on the surface of some types of cancer cells, including ovarian cancer. Your doctor may refer to it as a biomarker. Biomarkers can help determine if a targeted treatment is right for you



Before you start treatment, your doctor will test a sample from your tumor to confirm that your cancer is FR $\alpha$  positive. Your doctor may be able to use tissue from an earlier biopsy or surgery for testing



**ELAHERE is the first treatment** for people with platinum-resistant ovarian cancer that targets FR $\alpha$  in order to deliver medicine inside the cell



Biomarkers may be genes, proteins, or other materials in a cell that can be tested and measured to give more information about the cancer. This information can include how the tumor is growing, or how the cancer may respond to a type of treatment

FR $\alpha$ =folate receptor alpha.

## IMPORTANT SAFETY INFORMATION (CONT'D)


### What should I tell my healthcare provider before receiving ELAHERE?

**Tell your healthcare provider about all of your medical conditions, including if you:**

- have vision or eye problems.
- have numbness or tingling in your hands or feet.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**

# How does ELAHERE target and kill cancer cells?



## ELAHERE is the first targeted therapy developed and approved for FR $\alpha$ + platinum-resistant ovarian cancer

- ELAHERE is a type of treatment called an antibody-drug conjugate (ADC)
- ADCs are different from traditional chemotherapies and are designed to treat cancer in a targeted way

FR $\alpha$ =folate receptor alpha.

### IMPORTANT SAFETY INFORMATION (CONT'D)

- have liver problems.
- are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.


Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

# ELAHERE may help stop cancer from progressing



In a clinical study of 453 people with FR $\alpha$ -positive, platinum-resistant ovarian cancer, ELAHERE helped extend the time to disease progression compared with traditional chemotherapy.

## Median time to disease progression



**This is called median progression-free survival.**

Median progression-free survival is the length of time that the treatment stops the cancer from growing or getting worse for half of the people. A median is the middle number in a group of numbers that are listed from lowest to highest

**Traditional chemotherapy** in the clinical study included anti-cancer drugs that are not targeted and do not contain a material called platinum

FR $\alpha$ =folate receptor alpha.

## IMPORTANT SAFETY INFORMATION (CONT'D)

### Patients who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ELAHERE.
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.


Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

## Some people taking ELAHERE lived longer



ELAHERE was proven to help people live longer compared with traditional chemotherapy in a clinical study of 453 people with FR $\alpha$ -positive, platinum-resistant ovarian cancer.

### Median overall survival



#### **This is called median overall survival.**

Median overall survival is the length of time that half of the people were alive after starting treatment. A median is the middle number in a group of numbers that are listed from lowest to highest

FR $\alpha$ =folate receptor alpha.

### **IMPORTANT SAFETY INFORMATION (CONT'D)**

- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**




## A chance for better response is here



In the same clinical study, more people's tumors shrank by at least 30% or disappeared when treated with ELAHERE vs when treated with traditional chemotherapy.

### Overall response\*



The decrease in tumor size lasted a median of 6.8 months in those treated with ELAHERE and 4.5 months in those treated with traditional chemotherapy.



Overall response is a measure of all people who had either a complete or partial response

\*In the clinical study, 225 people treated with ELAHERE and 224 people treated with traditional chemotherapy were eligible to be evaluated for overall response

### IMPORTANT SAFETY INFORMATION (CONT'D)

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**

## What side effects may I expect on treatment?



### **ELAHERE can cause serious side effects, including:**

- **Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes
- **Lung problems (pneumonitis).** ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain
- **Peripheral neuropathy.** Nerve problems called peripheral neuropathy are common with ELAHERE and can also be severe. Your healthcare provider will monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if you get new or worsening numbness, tingling, burning sensation, or pain in your hands or feet or muscle weakness

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

## What side effects may I expect on treatment? (cont'd)



### The most common side effects and abnormal lab results include:

- Increased liver enzymes in the blood
- Feeling tired
- Blurred vision
- Nausea
- Diarrhea
- Stomach-area (abdominal) pain
- Changes in the cornea (part of the eye)
- Peripheral neuropathy
- Muscle, bone, or joint pain
- Decreased red or white blood cell counts
- Decreased platelets
- Decreased magnesium level in the blood
- Dry eye
- Constipation
- Vomiting
- Decreased albumin level in the blood
- Decreased appetite



Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have certain side effects

Due to side effects: 9% of people stopped treatment, 54% of people had dose delays, 34% of people had their dose reduced, and 24% of people reported serious adverse events

Of those treated with ELAHERE in a clinical study, 1% experienced hair thinning or hair loss.

**These are not all the possible side effects of ELAHERE.** Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).

Actor portrayal.



## How could ELAHERE affect my eyes?



People treated with ELAHERE can develop eye-related side effects, which can be severe. These may include blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes. Tell your doctor if you have a history of vision or eye problems.



Eye-related side effects most commonly occurred between the second and third infusion of ELAHERE



1% of people in the study stopped treatment because of eye problems

Your doctor will also have you meet with an eye doctor (optometrist or ophthalmologist) before your first infusion and periodically throughout your treatment. You will receive instructions for eye drops to use during your treatment. Be sure to use them as directed

Please refer to the ELAHERE Eye Care Brochure to learn about what you can do to help manage the risk of eye-related side effects during treatment with ELAHERE.

### IMPORTANT SAFETY INFORMATION (CONT'D)

**What are the possible side effects of ELAHERE? ELAHERE can cause serious side effects, including:**

- **Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**

## How should I care for my eyes?



### To help manage your risk of vision changes, here are some steps you can take:



- You should use steroid and lubricating eye drops as directed by your doctor. Your doctor will prescribe steroid eye drops and over-the-counter, lubricating, preservative-free eye drops for use before and during treatment with ELAHERE. Refer to the ELAHERE Eye Care Brochure for more detailed information
- Avoid wearing contact lenses throughout your treatment with ELAHERE unless your doctor tells you that you can
- **During treatment with ELAHERE, tell your doctor if you experience any eye problems, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes**

**Need reminders to use your eye drops?** There are several mobile apps available for download, designed to help you and/or your caregiver remember your eye drop schedule. You or your caregiver can find a variety to choose from at your preferred app store

### IMPORTANT SAFETY INFORMATION (CONT'D)

- **Lung problems (pneumonitis).** ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**

# When should I use my eye drops?




## You should use eye drops as directed by your doctor

### Steroid eye drops

Using steroid eye drops is recommended while taking ELAHERE.

#### Steroid Eye Drop Schedule for Each ELAHERE Infusion Cycle (Every 3 Weeks)



### Lubricating eye drops

The use of preservative-free lubricating eye drops is also recommended at least 4 times daily and as needed during treatment. **Wait at least 10 minutes after applying steroid eye drops** before using lubricating eye drops.

## Download the Eye Drop Tracker

While on treatment, use this tool to help track your eye drop use, notes to share with your doctor, and upcoming appointments

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).



Actor portrayals.

## What happens before I receive an ELAHERE infusion?



Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking ELAHERE with certain other medicines may cause side effects.

- Before each dose of ELAHERE, you will receive medicines to help prevent infusion-related reactions, nausea, and vomiting
  - Some of these medications may be taken orally and some may be given to you by infusion into your vein
  - Usually, these medications will be given on the same day as the infusion. However, your doctor may choose to give you the medications the day before you start ELAHERE

**REMINDER:** Your doctor will test a tissue sample to confirm your cancer is FR $\alpha$  positive to help determine if ELAHERE may be right for you. To do this, they can use tumor tissue stored from an earlier biopsy or surgery or take a new sample

FR $\alpha$ =folate receptor alpha.

### IMPORTANT SAFETY INFORMATION (CONT'D)

- **Peripheral neuropathy.** Nerve problems called peripheral neuropathy are common during treatment with ELAHERE and can also be severe. Your healthcare provider will monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if you get new or worsening numbness, tingling, burning sensation or pain in your hands or feet or muscle weakness.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

## What should I discuss with my doctor before my first infusion?



### **Before receiving ELAHERE, tell your doctor about all of your medical conditions, including if you:**

- Have vision or eye problems
- Have numbness or tingling in your hands or feet
- Have liver problems
- Are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your doctor right away if you become pregnant or think you may be pregnant during treatment with ELAHERE
- Are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE

### **If you are able to become pregnant:**

- Your doctor should do a pregnancy test before you start treatment with ELAHERE
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE

### **IMPORTANT SAFETY INFORMATION (CONT'D)**


**The most common side effects and abnormal labs of ELAHERE include:** • increased liver enzymes in the blood • feeling tired • blurred vision • nausea • diarrhea • stomach-area (abdominal) pain • changes in the cornea (part of the eye) • peripheral neuropathy • muscle, bone, or joint pain • decreased red or white blood cell counts • decreased platelets • decreased magnesium level in the blood • dry eye • constipation • vomiting • decreased albumin level in the blood • decreased appetite

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including [Boxed WARNING](#) and [Medication Guide](#).**

## How will I receive ELAHERE?



**ELAHERE is given by infusion (IV or a port) every 3 weeks**



ELAHERE dosage is determined by your weight. To ensure accurate dosing, your doctor will weigh you prior to treatment.

If you have side effects with ELAHERE, your doctor may reduce your dose, delay your dose, or completely stop treatment

IV=intravenous.

### **IMPORTANT SAFETY INFORMATION (CONT'D)**

Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have certain side effects.

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).**



# What should I ask my doctor?



It's important to ask questions and start a conversation about what is next in your treatment journey. The below list of questions can help guide your conversation with your doctor and care team.

- What is FR $\alpha$  and why is it important?
- Have I been tested for FR $\alpha$ , the protein that ELAHERE targets? What were the results?
- What is platinum resistance?
- How is ELAHERE different from other treatments I have been on?
- What is the goal of treatment with ELAHERE?
- What are the side effects I might experience with this treatment?
- Is there anything I can do to manage those side effects myself?
- Will I need to see other specialized doctors?
- What can I do to get ready for treatment?
- Should I go to my regular eye doctor?
- **Question for your eye doctor:** My oncologist put me on a drug that can affect my eyes. Would your office be able to work with my oncologist to help monitor my eye health during my treatment?

## ELAHERE Starter Kit and Welcome Kit

Ask your doctor about the ELAHERE Starter Kit and sign up to receive a Welcome Kit. These kits include resources to help support you when beginning treatment with ELAHERE

FR $\alpha$ =folate receptor alpha.

Please see Important Safety Information throughout and accompanying [Full Prescribing Information](#), including **Boxed WARNING** and [Medication Guide](#).

Actor portrayals.



# Is there financial support for ELAHERE?



## ELAHERE Support Services: Here to help you get started on ELAHERE

Once enrolled in the program, your ELAHERE Support Services Case Manager will help you understand your insurance coverage and navigate financial assistance options.

\$0

### For patients with commercial insurance\*:

Eligible patients could pay  
as little as \$0 for their medication



### Patient Assistance Program (PAP)

The Patient Assistance Program (PAP) provides eligible patients with ELAHERE at no charge<sup>†</sup>

\*Terms and conditions apply. Patients are eligible for co-pay assistance if enrolled in private commercial health insurance and are not covered by state or federal healthcare programs, and who meet the eligibility criteria. Patients will be enrolled for 12 months. There are no income requirements to participate in the program.

<sup>†</sup>Criteria include: patients who are uninsured or have insurance that excludes coverage for ELAHERE (including patients on Medicare or Medicaid), residents of the United States or Puerto Rico, and patients who meet the financial eligibility requirements. Terms and conditions apply.

## IMPORTANT SAFETY INFORMATION (CONT'D)

These are not all of the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit

[www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including [Boxed WARNING](#) and [Medication Guide](#).

Actor portrayal.

Here for

More



**We're here to help. Contact us to learn about our support programs and financial assistance.**

For questions, call **1-833-ELAHERE**  
Monday-Friday, 8:00 AM to 8:00 PM ET  
Or email **[ELAHESupport@cardinalhealth.com](mailto:ELAHESupport@cardinalhealth.com)**

Visit **[ELAHERE.com](https://www.elaHERE.com)** to learn more  
and find additional resources

## SELECT IMPORTANT SAFETY INFORMATION

**What is the most important information I should know about ELAHERE?**

**ELAHERE can cause serious side effects, including:**

**Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes. **Continue reading information about eye problems on pages 3 and 5.**

**Please see additional Important Safety Information throughout and accompanying [Full Prescribing Information](#), including Boxed WARNING and [Medication Guide](#).**



© 2024 ImmunoGen, Inc., All Rights Reserved. US-MIRV-2300291 03/2024  
ImmunoGen, ELAHERE, and their respective logos are trademarks or registered trademarks of ImmunoGen, Inc.